

# Anisotropic static solutions in modelling highly compact bodies

M. Chaisi\* and S. D. Maharaj<sup>†</sup>

School of Mathematical Sciences

University of KwaZulu-Natal

Durban 4041

South Africa.

## Abstract

Einstein field equations for anisotropic spheres are solved and exact interior solutions obtained. This paper extends earlier treatments to include anisotropic models which accommodate a wider variety of physically viable energy densities. Two classes of solutions are possible. The first class contains the limiting case  $\mu \propto r^{-2}$  for the energy density which arises in many astrophysical applications. In the second class the singularity at the center of the star is not present in the energy density. The models presented in this paper allow for increasing and decreasing profiles in the behavior of the energy density.

---

\*Permanent address: Department of Mathematics & Computer Science, National University of Lesotho, Roma 180, Lesotho.; eMail: [m.chaisi@nul.ls](mailto:m.chaisi@nul.ls)

<sup>†</sup>eMail: [maharaj@ukzn.ac.za](mailto:maharaj@ukzn.ac.za)

# 1 Introduction

Anisotropic spheres in general relativity, in spherically symmetric spacetimes which are not static, have continued to attract the attention of researchers since the pioneering work of Bowers and Liang<sup>1</sup>. Some of the interesting recent papers published in this regard include the works of Dev and Gleiser<sup>2,3</sup>, Herrera *et al*<sup>4</sup>, Ivanov<sup>5</sup>, Mak and Harko<sup>6</sup>, and Sharma and Mukherjee<sup>7</sup>. The original reason for studying relativistic anisotropic spheres was to generate models that permit redshifts higher than the critical redshift  $z_c$  of isotropic matter. The analysis of Bondi<sup>8</sup> demonstrated the feasibility and stability of higher redshifts for anisotropic Newtonian and relativistic stars. Anisotropy cannot be neglected in stellar clusters and galaxies, in addition to individual stars, as pointed out by Binney and Tremaine<sup>9</sup>, Cuddeford<sup>10</sup> and Michie<sup>11</sup>. Anisotropy may be an intrinsic feature of boson stars as suggested by Dev and Gleiser<sup>2</sup> and Mak and Harko<sup>6</sup>. Sharma and Mukherjee<sup>7</sup> considered the theoretical possibility of anisotropy in strange stars, with densities greater than neutron stars but less than black holes, which are bound stable stellar bodies.

In a recent treatment Chaisi and Maharaj<sup>12</sup> found an exact anisotropic solution to the field equations utilizing an algorithm of Maharaj and Maartens<sup>13</sup>. This solution has a simple form: it can be expressed completely in terms of elementary functions and it has a clear physical interpretation. In this analysis we extend the Chaisi and Maharaj<sup>12</sup> solution to include anisotropic models which allow for a wide variety of densities which are physically reasonable. We present four classes of solution and discuss their physical features. In §2 we present the relevant classes of solutions in the form of tables to ensure clarity and to highlight the interconnections between the classes. In §3 the physical features of our solutions are briefly discussed, and §4 contains the discussion.

## 2 New classes of solutions

The Einstein field equations to be integrated are

$$e^{-\lambda} = 1 - \frac{2m}{r} \quad (1)$$

$$r(r - 2m)\nu' = p_r r^3 + 2m \quad (2)$$

$$(\mu + p_r)\nu' + 2p_r' = -\frac{4}{r}(p_r - p_\perp) \quad (3)$$

In the above we have defined  $m(r) = \frac{1}{2} \int_0^r x^2 \mu(x) dx$  which is the mass function. The radial pressure  $p_r = p + 2S/\sqrt{3}$  and the tangential pressure  $p_\perp = p - S/\sqrt{3}$  are not equal for anisotropic matter. The magnitude  $S$  provides a measure of anisotropy.

The field equations (1)-(3) were integrated by Chaisi and Maharaj<sup>12</sup> for the energy density

$$\mu = \frac{j}{r^2} + k + \ell r^2 \quad (4)$$

where  $j, k, \ell$  are constants. It is possible to integrate the field equations for other choices of the energy density which are physically acceptable. In this section we present three new classes of exact solutions to supplement the class found earlier.

| Case | $\mu(r)$                                                                                              |
|------|-------------------------------------------------------------------------------------------------------|
| I    | $\frac{j}{r^2} + k + \ell r^2$                                                                        |
| II   | $\frac{1}{r^2} (1 - a) - (p + 1)br^{p-2}$                                                             |
| III  | $\frac{1}{r^2} \left( \frac{a-c+(b-d)r^2}{a+br^2} \right) - 2 \frac{d(a+br^2)-b(c+dr^2)}{(a+br^2)^2}$ |
| IV   | $\frac{1}{r^2} \left( \frac{a-1+br^2+cr^4}{a+br^2+cr^4} \right) + 2 \frac{b+2cr^2}{(a+br^2+cr^4)^2}$  |

Table 1: Energy density functions

Table 1 comprises a list of forms for energy densities that have been studied. The forms of energy density  $\mu$  in Table 1 were chosen so that the Einstein field equations could be fully integrated, and the gravitational potentials and matter variables written in closed form. The functions chosen for  $\mu$  in Table 1 have profiles which correspond to physically acceptable anisotropic spheres, and their simple forms facilitate the analysis of the gravitational potentials and the matter variables. To complete the integration of the Einstein

field equations we also need to make a choice for the radial pressure  $p_r$ . Clearly a variety of choices for  $p_r$  is possible; our choice is made on physical grounds. The following is a list of forms for the radial pressure:

| Case | $p_r(r)$                                                                                                   |
|------|------------------------------------------------------------------------------------------------------------|
| I    | $\frac{C}{1-j} \left(1 - j - \frac{k}{3}r^2 - \frac{\ell}{5}r^4\right) \left(1 - \frac{r^2}{R^2}\right)^n$ |
| II   | $\frac{C}{a} (a + br^p) \left(1 - \frac{r^2}{R^2}\right)^n$                                                |
| III  | $\frac{a}{c} C \left(\frac{c+dr^2}{a+br^2}\right) \left(1 - \frac{r^2}{R^2}\right)^n$                      |
| IV   | $\frac{aC}{a+br^2+cr^4} \left(1 - \frac{r^2}{R^2}\right)^n$                                                |

Table 2: Radial pressure functions

The forms of  $p_r$  selected in Table 2 all reduce to the expression  $p_r = C(1 - r^2/R^2)^n$  with appropriate choices for the parameters so that the radial pressure is a monotonically decreasing function. Table 3 contains the gravitational potentials  $e^\nu$  and  $e^\lambda$ . Table 4 lists the corresponding matter variables: the energy density  $\mu$ , the mass function  $m$ , the radial pressure  $p_r$  and the tangential pressure  $p_\perp$ , respectively.

We believe that the families of solutions for Cases I-IV presented in Tables 3-4 are new solutions to the field equations, apart from particular special cases, for relativistic anisotropic matter. Case I contains the Maharaj and Maartens<sup>13</sup> solution ( $j = \ell = 0$ ) and the Gokhroo and Mehra<sup>14</sup> solution ( $j = 0$ );  $I_2$  is defined in terms of elementary functions<sup>12</sup>. The exact solutions in Cases I-IV are amenable to a physical analysis because they have a simple form, and in all cases the gravitational potentials and matter variables are given in terms of elementary functions. The parameters  $j$ ,  $k$ , and  $\ell$  in Case I are constants. The quantities  $a$ ,  $b$ ,  $c$ ,  $d$  and  $p$  are also constants in Cases II-IV. Note that the constant  $C$  corresponds to central radial pressure ( $C = p_r(0), n \geq 1$ )

| Case | $e^\lambda$                                                    | $e^\nu$                                                                                                                                 |
|------|----------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------|
| I    | $\left(1 - j - \frac{k}{3}r^2 - \frac{\ell}{5}r^4\right)^{-1}$ | $Br^{\frac{j}{1-j}} \exp \left\{ \frac{I_2}{1-j} - \frac{CR^2}{2(1-j)(n+1)} \left(1 - \frac{r^2}{R^2}\right)^{n+1} \right\}$            |
| II   | $(a + br^p)^{-1}$<br>$(p > 0)$                                 | $\frac{Br^{\frac{1-a}{a}}}{(a+br^p)^{\frac{1}{ap}}} \exp \left\{ -\frac{CR^2}{2a(n+1)} \left(1 - \frac{r^2}{R^2}\right)^{n+1} \right\}$ |
| III  | $\frac{a+br^2}{c+dr^2}$                                        | $Br^{\frac{a-c}{c}} (c + dr^2)^{\frac{bc-ad}{2cd}} \exp \left\{ -\frac{aCR^2}{2c(n+1)} \left(1 - \frac{r^2}{R^2}\right)^{n+1} \right\}$ |
| IV   | $a + br^2 + cr^4$                                              | $Br^{a-1} \exp \left\{ -\frac{aCR^2}{2(n+1)} \left(1 - \frac{r^2}{R^2}\right)^{n+1} + \frac{br^2}{2} + \frac{cr^4}{4} \right\}$         |

Table 3: Gravitational potentials

### 3 Physical features

The gravitational potentials  $e^\lambda$  are finite for all Cases I-IV at the centre  $r = 0$  and at the boundary  $r = R$ . The functions  $e^\lambda$  are continuous and well behaved in the interior of the relativistic star. The gravitational potentials  $e^\nu$  for all Cases I-IV are continuous and well behaved in the interior and finite at the boundary of the star  $r = R$ . However we observe from Table 3 that there is a singularity at the centre  $r = 0$  in general for all Cases I-IV in the potential  $e^\nu$ . The singularity in  $e^\nu$  is not present for specific choices of parameter values and may be removed by setting

$$j = 0, \quad \text{in Case I,} \quad (5)$$

$$a = 1, \quad \text{in Cases II and IV,} \quad (6)$$

$$a = c, \quad \text{in Case III.} \quad (7)$$

The gravitational potentials in Table 3 have the advantage of having a simpler analytic form, and they are written in terms of polynomials, rational and exponential functions. Consequently the radial and tangential pressures have a simple analytic representation.

The radial pressure  $p_r$  is continuous and well behaved in the interior. Also  $p_r > 0$  in the interval  $(0, R)$ , regular at the centre ( $p_r(r = 0) = C$ ), and vanishes at the boundary ( $p_r(r = R) = 0$ ) in all four cases. The tangential pressure  $p_\perp$  in the four cases has a singularity at the centre, but is otherwise well behaved throughout the interior of the star and finite at the boundary. Note that the singularity in  $p_\perp$  may be avoided by suitable

particular choices for parameter values. In general the tangential pressure is not zero at the boundary of the star ( $p_{\perp}(r = R) \neq 0$ ) and does not vanish, as does the radial pressure ( $p_r(r = R) = 0$ ). It is also important to observe that the magnitude of the stress tensor  $S$  is a nonzero function in general for all Cases I-IV. Hence this class of solutions is generally anisotropic and it is not possible for  $S$  to vanish and obtain an isotropic limit.

The energy density  $\mu$  for all cases contains the limiting case

$$\mu \propto r^{-2} \quad (8)$$

as can be verified directly from Table 4. It has been demonstrated that the energy density (8) arises in isothermal spheres for isotropic matter for both Newtonian and relativistic stars by Saslaw *et al*<sup>15</sup>. There are many other examples where the form (8) arises in relativistic astrophysics. Misner and Zapolsky<sup>16</sup> proposed that isotropic solutions (with  $k = 0 = \ell$ ) model the physical configuration of a relativistic Fermi gas for some particular value of the parameter  $j$ . Another example, given by Dev and Gleiser<sup>2</sup> who indicate that for particular values of  $j \neq 0$  and  $k \neq 0$  ( $\ell = 0$ ) the energy density  $\mu$  is appropriate for modelling the relativistic Fermi gas core immersed in a constant density background. Consequently our exact solutions in Cases I-IV, all containing the limiting energy density (8), may be used to model a variety of similar astrophysical situations when the anisotropy is not negligible.

It is also important to observe that our approach allows us to consider energy densities for which (8) does not hold. For this scenario we must utilize the values for the constants in (5)-(7). With the help of (5)-(7) we observe from Table 4 that the singularity at  $r = 0$  is removed and  $\mu$  is a continuous function throughout the interval  $[0, R]$ . Figure 1 provides an illustration of the behaviour of the energy density  $\mu$ , when (5)-(7) is true, for particular chosen values of the constants. The radial distance is over the interval  $0 \leq r \leq 1$ . From Figure 1 we observe that  $\mu$  is a well behaved function in the interior of the star and has finite values at the center and the boundary. Cases I-IV admit both possibilities of  $\mu' < 0$  and  $\mu' > 0$ ; that is anisotropic stars with decreasing or increasing energy densities from the centre to the boundary, respectively, can be studied.

| Case | $\mu$                                                                                                 | $m$                                                                   | $p_r$ ( $C = p_r(0), n \geq 1$ )                                                                                        | $p_\perp$                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
|------|-------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| I    | $\frac{j}{r^2} + k + \ell r^2$                                                                        | $\frac{r}{2} \left( j + \frac{k}{3} r^2 + \frac{\ell}{5} r^4 \right)$ | $\frac{C}{1-j} \left( 1 - j - \frac{k}{3} r^2 - \frac{\ell}{5} r^4 \right) \times \left( 1 - \frac{r^2}{R^2} \right)^n$ | $p_r + \frac{C}{2(1-j)} \left( j - \frac{\ell}{5} r^4 \right) \left( 1 - \frac{r^2}{R^2} \right)^n + \frac{r^2}{2} \left( 1 - j - \frac{k}{3} r^2 - \frac{\ell}{5} r^4 \right)^{-1} \times \left\{ \frac{C^2}{2(1-j)^2} \left( 1 - j - \frac{k}{3} r^2 - \frac{\ell}{5} r^4 \right)^2 \left( 1 - \frac{r^2}{R^2} \right)^{2n} - \frac{2nC}{(1-j)R^2} \left( 1 - j - \frac{k}{3} r^2 - \frac{\ell}{5} r^4 \right)^2 \left( 1 - \frac{r^2}{R^2} \right)^{n-1} + \frac{1}{2r^2} \left( \frac{j}{r^2} + k + \ell r^2 \right) \left( j + \frac{k}{3} r^2 + \frac{\ell}{5} r^4 \right) \right\}$ |
| II   | $\frac{1}{r^2} (1 - a)$<br>$-(p+1)br^{p-2}$                                                           | $\frac{r}{2} (1 - a - br^p)$<br>$(p > 0)$                             | $\frac{C}{a} (a + br^p) \times \left( 1 - \frac{r^2}{R^2} \right)^n$                                                    | $p_r + \frac{C}{a} \left( \frac{1-a}{2} + b \left( \frac{p}{4} - \frac{1}{2} \right) r^p \right) \left( 1 - \frac{r^2}{R^2} \right)^n + \frac{r^2}{2} (a + br^p)^{-1} \times \left\{ \frac{C^2}{2a^2} (a + br^p)^2 \left( 1 - \frac{r^2}{R^2} \right)^{2n} - \frac{2nC}{aR^2} (a + br^p)^2 \left( 1 - \frac{r^2}{R^2} \right)^{n-1} + \frac{1}{2r^4} (1 - a - br^p) (1 - a - b(p+1)r^p) \right\}$                                                                                                                                                                                          |
| III  | $\frac{1}{r^2} \left( \frac{a-c+(b-d)r^2}{a+br^2} \right) - \frac{2(ad-bc)}{(a+br^2)^2}$              | $\frac{r}{2} \left( 1 - \frac{c+dr^2}{a+br^2} \right)$                | $\frac{a}{c} C \left( \frac{c+dr^2}{a+br^2} \right) \times \left( 1 - \frac{r^2}{R^2} \right)^n$                        | $p_r + \frac{r}{4} \left\{ \frac{a-c+(b-d)r^2}{r^2(a+br^2)} - \frac{2(ad-bc)}{(a+br^2)^2} + \frac{aC}{c} \left( \frac{c+dr^2}{a+br^2} \right) \left( 1 - \frac{r^2}{R^2} \right)^n \right\} \times \left\{ \frac{aCr}{c} \left( 1 - \frac{r^2}{R^2} \right)^n + \frac{a-c+(b-d)r^2}{r(c+dr^2)} \right\} + \frac{aCr^2}{c} \frac{ad-bc}{(a+br^2)^2} \left( 1 - \frac{r^2}{R^2} \right)^n - \frac{anCr^2}{cR^2} \left( \frac{c+dr^2}{a+br^2} \right) \left( 1 - \frac{r^2}{R^2} \right)^{n-1}$                                                                                               |
| IV   | $\frac{1}{r^2} \left( \frac{a-1+br^2+cr^4}{a+br^2+cr^4} \right) + \frac{2(b+2cr^2)}{(a+br^2+cr^4)^2}$ | $\frac{r}{2} \left( 1 - \frac{1}{a+br^2+cr^4} \right)$                | $\frac{aC}{a+br^2+cr^4} \times \left( 1 - \frac{r^2}{R^2} \right)^n$                                                    | $p_r + \frac{r}{4} \left\{ \frac{a-1+br^2+cr^4}{r^2(a+br^2+cr^4)} + \frac{2(b+2cr^2)}{(a+br^2+cr^4)^2} + \frac{aC}{a+br^2+cr^4} \left( 1 - \frac{r^2}{R^2} \right)^n \right\} \times \left\{ aCr \left( 1 - \frac{r^2}{R^2} \right)^n + \frac{1}{r} (a - 1 + br^2 + cr^4) \right\} - aCr^2 \frac{b+2cr^2}{(a+br^2+cr^4)^2} \left( 1 - \frac{r^2}{R^2} \right)^n - \frac{aCnr^2}{R^2(a+br^2+cr^4)} \left( 1 - \frac{r^2}{R^2} \right)^{n-1}$                                                                                                                                                |

Table 4: Matter variables

## 4 Discussion

We have presented four new classes of solution which model anisotropic stars. These solutions may be broadly divided into two categories in terms of the behaviour of the energy density  $\mu$  at the centre  $r = 0$ : the first category contains the limiting case  $\mu \propto r^{-2}$ , and in the second category the singularity at the centre of the star is not present. The solutions found permit  $\mu' < 0$  and  $\mu' > 0$  which allow for decreasing and increasing energy densities as we move from the centre to the boundary of the star. Note that the solutions presented in this paper have the feature that  $p_r \neq p_\perp$  in general so that the anisotropy factor  $S \neq 0$ ; consequently our solutions do not have an isotropic limit.

We make three observations relating to the physical reasonableness of our model. Firstly, the vanishing of the pressure at the boundary is a consequence of the first and second fundamental forms; there is no restriction placed on the tangential pressure which may be nonvanishing. This feature is also evident in the solutions of Chaisi and Maharaj<sup>12</sup>, Maharaj and Maartens<sup>13</sup> and Gokhroo and Mehra<sup>14</sup>. Secondly, the solutions in this paper do not satisfy an equation of state. For a realistic astrophysical matter distribution we should impose a barotropic equation of state relating the pressure and energy density; this will be pursued in future work. Thirdly, we observe that for realistic models the space derivatives of the matter field should be negative. This will have the effect of restricting the arbitrary constants in the energy density; for example this would place restrictions on the relative magnitudes of  $j, k, \ell$  of Case I.




Figure 1: Energy density  $\mu$  plots with singularities removed



# Acknowledgements

SDM and MC thank the National Research Foundation of South Africa for financial support. MC is grateful to the University of KwaZulu-Natal for a scholarship. We are thankful to the referee for insightful comments.

# References

- [1] R.L. Bowers and E.P.T. Liang, *Astrophys. J.*, **188**, 657, (1974)
- [2] K. Dev and M. Gleiser, *Gen. Rel. Grav.*, **34**, 1793, (2002)
- [3] K. Dev and M. Gleiser, *Gen. Rel. Grav.*, **35**, 1435, (2003)
- [4] L. Herrera and J. Martin and J. Ospino, *J. Math. Phys.*, **43**, 4889, (2002)
- [5] B.V. Ivanov, *Phys. Rev. D*, **65**, 10411, (2002)
- [6] M.K. Mak and T. Harko, *Chin. J. Astron. Astrophys.*, **2**, 248, (2002)
- [7] R. Sharma and S. Mukherjee, *Mod. Phys. Lett. A*, **17**, 2535, (2002)
- [8] H. Bondi, *Mon. Not. R. Astron. Soc.*, **259**, 365, (1992)
- [9] J. Binney and S. Tremaine, *Galactic Dynamics*, Princeton University Press, Princeton, NJ, (1987)
- [10] P. Cuddeford, *Mon. Not. R. Astr. Soc.*, **253**, 414, (1991)
- [11] R.W. Michie, *Mon. Not. R. Astr. Soc.*, **125**, 127, (1963)
- [12] M. Chaisi and S.D. Maharaj, *Gen. Rel. Grav.*, **37** 1177 (2005)
- [13] S.D. Maharaj and R. Maartens, *Gen. Rel. Grav.*, **21**, 899 (1989)
- [14] M.K. Gokhroo and A.L. Mehra, *Gen. Rel. Grav.*, **26**, 75, (1994)
- [15] W.C. Saslaw and S.D. Maharaj and N Dadhich, *Astrophys. J.*, **471**, 571, (1996)
- [16] C.W. Misner and H.S. Zepolsky, *Phys. Rev. Lett.*, **12**, 635, (1964)